

# Barricade® 0.38% Plus Fertilizer

For selective preemergence control of grass and broadleaf weeds in established turfgrasses (excluding golf course putting greens), lawns, athletic fields, parks, and sod farms, field grown and landscape ornamentals, established perennials and wildflower plantings, Non-crop areas, including landscape beds, ornamentals, and established perennial wildflower plantings on or surrounding municipal grounds and government installations.

**ACTIVE INGREDIENT:**  
Prodiamine: (N<sup>3</sup>, N<sup>3</sup>-Di-n-propyl-2,4-dinitro-6-(trifluoromethyl)-m-phenylenediamine ..... 0.38%  
**OTHER INGREDIENTS:**..... 99.62%  
Total ..... 100.00%

KEEP OUT OF REACH OF CHILDREN

CAUTION

See the following sections Precautionary Statements, Hazards to Humans and Domestic Animals, for additional information.

FIRST AID

If on skin or clothing	<ul style="list-style-type: none"><li>Take off contaminated clothing.</li><li>Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>Call a poison control center or doctor for treatment advice.</li></ul>
If in eyes	<ul style="list-style-type: none"><li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li><li>Call a poison control center or doctor for treatment advice.</li></ul>
If inhaled	<ul style="list-style-type: none"><li>Move person to fresh air.</li><li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to mouth if possible.</li><li>Call a poison control center or doctor for treatment advice.</li></ul>

HOT LINE NUMBER

For 24 Hour Medical Emergency Assistance ( Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire, or Accident), call CHEM-TREC 1-800-424-9300

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-424-9300 for emergency medical treatment information.

**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**  
**CAUTION:** Harmful if absorbed through the skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. [Wear protective eyewear]. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Wear long sleeved shirt and long pants, shoes plus socks, and chemical-resistant waterproof gloves. Remove and wash contaminated clothing before reuse. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.  
**Personal Protective Equipment (PPE)**  
**Mixers, loaders, applicators, and other handlers must wear:**

- Long-sleeved shirt and long pants
- Shoes plus socks
- Chemical-resistant gloves made of any water proof material

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

**ENGINEERING CONTROL STATEMENTS**  
When handlers used closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240 (d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.  
**User Safety Recommendations:**  
**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing /PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**Environmental Hazards**  
This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment wash waters.  
**DIRECTIONS FOR USE**  
It is a violation of federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the State/Tribal agency responsible for pesticide regulation.

- For use on plants intended for aesthetic purposes or climatic modification.
- Read entire label before using this product.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.  
**Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.** Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.  
PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks


NON AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.  
**Do not enter or allow others to enter until dusts have settled.**

**ESTABLISHED TURF**  
Golf courses (except putting greens), lawns, parks, athletic fields, sod farms.  
**How to Apply**

- Do not apply more than 9.0 lbs. product/1,000 sq. ft. (1.5 lb. ai/A) per calendar year.
- Do not make more than two applications per calendar year.
- Refer to Rates for Grass Species in table and Granular Spreader Settings table for spreader settings to deliver the rate recommended for your lawn grass species. Uniform coverage is necessary to provide the most effective level of control.
- Apply LESCO Barricade 0.38% Plus Fertilizer to the turf in the early spring prior to crabgrass emergence. See Crabgrass Germination Dates in the map below for the approximate timing in your area. Always apply before weeds emerge. LESCO Barricade 0.38% Plus Fertilizer will not control emerged crabgrass.
- If you have a known serious weed problem or live in an area with an extended growing season, you may need to make a second application 60-90 days later.
- If you live in Zones D and E (see Crabgrass Germination Dates map), you may apply a higher rate (Medium or High Rate) or make two applications for more effective crabgrass control, if recommended for your turfgrass species.
- Do not make a second application to Fine Fescue, Creeping Bentgrass, and newly sprigged and plugged Bermudagrass. Do not make a second application if you applied the High Rate to Buffalograss, Kentucky Bluegrass, or Perennial Ryegrass.

For best control – After application, water in ½ inch of water within 14 days, either from watering or rain.



Turf Species	Application Schedule <sup>1</sup>	Application Rate <sup>2</sup>		
		Base	Medium	High
Bermudagrass (established) Bahia Centipedegrass Kikuyugrass Seashore Paspalum St. Augustinegrass Tall Fescue (including turf-type) Zoysiagrass	First Application			
	Second Application (optional)	Base or Medium or High	Base or Medium or High	Base or Medium or High
Buffalograss Kentucky Bluegrass Perennial Ryegrass	First Application	Base	Medium	High
	Second Application (optional)	Base or Medium	Base or Medium	-
Fine Fescue (including Red Fescue)	One Application Only	Base or Medium or High	-	-
Creeping Bentgrass (½ inch and greater height) <sup>3</sup>	One Application Only	Base or Medium	-	-
Bermudagrass newly sprigged and plugged) <sup>4</sup>	One Application Only	Base or Medium	-	-

- A second application may be necessary because of heavy weed pressure or extended growing season.
- See Ggranular Spreader Settings table for spreader settings to deliver the rate.
- To avoid grass injury, do not apply LESCO Barricade 0.38% Plus Fertilizer to creeping bentgrass mowed at less than ½ inch in height.
- Newly sprigged or plugged Bermudagrass rooting may be temporarily slowed.

13-0-4

GUARANTEED ANALYSIS

Total Nitrogen (N) ..... 13.00%  
13.00% Urea Nitrogen \*  
0.00% Water Insoluble Nitrogen  
0.00% Other Water Soluble Nitrogen

Soluble Potash (K<sub>2</sub>O) ..... 4.00%

Sulfur (S) Total ..... 1.00%  
1.00% Free Sulfur (S)

DERIVED FROM: Polymer Coated Sulfur Coated Urea, Urea, Muriate of Potash.

CHLORINE (Cl) MAX. .... 3.00%

\*2.92% Slowly available Urea Nitrogen from Polymer Coated Sulfur Coated Urea.

#097649  
T

# Barricade® 0.38% Plus Fertilizer

## TOLERANT ORNAMENTALS

LESCO Barricade 0.38% Plus Fertilizer will not harm most trees, shrubs, vines and flowers. See the table below: Table 1. Tolerant Ornamentals – All States

Table 1. Tolerant Ornamentals – All States

Scientific Name	Common Name	Scientific Name	Common Name
<i>Abies</i> spp.	Fir Species (Balsam, Fraser, Noble, etc.)	<i>Maleophora luteola</i>	Ice Plant
<i>Acer palmatum</i>	Japanese Maple	<i>Malus</i> sp.	Crabapple*
<i>Acer platanoides</i>	Norway Maple	<i>Nandina domestica</i>	Heavenly Bamboo
<i>Agapanthus africanus</i>	Lily-of-the-Nile	<i>Narcissus</i> spp.	Narcissus species
<i>Arctotheca calendula</i>	Cape Weed	<i>Nerium</i> spp.	Oleander
<i>Aucuba japonica</i>	Japanese Aucuba	<i>Olea europaea</i>	Olive*
<i>Berberis gladywynensis</i>	Barberry	<i>Ophiopogon japonicus</i>	Mondo Grass
<i>Berberis julianae</i>	Wintergreen Barberry	<i>Osteospermum fruticosum</i>	Trailing African Daisy
<i>Berberis thunbergii</i>	Japanese Barberry	<i>Oxydendrum arboreum</i>	Sourwood
<i>Berberis verruculosa</i>	Warty Barberry	<i>Persea americana</i>	Avocado*
<i>Buxus microphylla</i>	Japanese Boxwood	<i>Photinia fraseri</i>	Fraser's Photinia (Redtip)
<i>Callistemon viminalis</i>	Weeping Bottlebrush	<i>Picea</i> spp.	Colorado Blue, Norway, etc.
<i>Calluna vulgaris</i>	Scotch Heather	<i>Pieris japonica</i>	Lily-of-the-Valley Shrub
<i>Carpobrotus edulis</i>	Hottentot Fig (Ice Plant)	<i>Pinus elliotii</i>	Slash Pine
<i>Cassia artemisoides</i>	Feathery Cassia	<i>Pinus nigra</i>	Austrian Black Pine
<i>Ceanothus rigidus</i>	Wild Lilac	<i>Pinus radiata</i>	Monterey Pine
<i>Chamaecyparis pisifera</i>	False Cypress	<i>Pinus palustris</i>	Longleaf Pine
<i>Cleyera japonica</i>	Cleyera	<i>Pinus strobus</i>	Eastern White Pine
<i>Citrus</i> spp.	Citrus Species*	<i>Pinus sylvestris</i>	Scotch Pine
<i>Cornus florida</i>	Flowering Dogwood	<i>Pinus taeda</i>	Loblolly Pine
<i>Cornus stolonifera</i>	American Dogwood	<i>Pinus thunbergiana</i>	Japanese Black Pine
<i>Cortaderia selloana</i>	Pampas Grass	<i>Pinus virginiana</i>	Virginia Pine
<i>Cotoneaster apiculatus</i>	Cranberry Cotoneaster	<i>Pistachio</i> sp.	Pistachio*
<i>Cotoneaster buxifolius</i>	Cotoneaster	<i>Pittosporum rhombifolium</i>	Queensland Pittosporum
<i>Cotoneaster microphyllus</i>	Rockspray Cotoneaster	<i>Pittosporum tobira</i>	Japanese Pittosporum
<i>Cretaeagus</i> spp.	Hawthorne	<i>Prunus laurocerasus</i>	English Laurel
<i>Cupressus sempervirens</i>	Italian Cypress	<i>Prunus</i> spp.	Almond, Apricot, Nectarine, Peach, Plum, and Prune*
<i>Delasperma alba</i>	White Trailing Ice Plant	<i>Pseudotsuga menziesii</i>	Douglas Fir
<i>Dodonea viscosa</i>	Hop Bush	<i>Pyracantha fortuneana</i> ; <i>koidzumii</i>	Firethorn spp.
<i>Elaeagnus pungens</i>	Silverberry	<i>Pyrus</i> sp.	Bradford Pear sp.
<i>Euonymus fortunei</i>	Wintercreeper	<i>Quercus</i> spp.	Oak spp.
<i>Euonymus japonica</i>	Evergreen Euonymus	<i>Raphiolepis indica</i>	Indian Hawthorne
<i>Euonymus kiautschovica</i>	Spreading Euonymus		
<i>Fatsia japonica</i>	Japanese Aralia	<i>Rhododendron (including azalea)</i>	'Coral Bells' 'Formosa', 'Hino-crimson', 'PJM', 'Roseum Elegans'
<i>Forsythia intermedia</i>	Border Forsythia	<i>Rosmarinus officinalis</i>	Rosemary
<i>Gardenia jasminoides</i>	Gardenia, Cape-Jasmine	<i>Rumohra adiantiformis</i>	Leathleaf Fern
<i>Gladiolus</i> spp.	Gladiolus species	<i>Santolina virens</i>	Santolina
<i>Hedera helix</i>	English Ivy	<i>Sedum album</i>	Stonecrop
<i>Hibiscus</i>	Rose of Sharon	<i>Syzygium paniculatum</i>	Japanese Boxcherry
<i>Ilex cornuta</i>	Chinese Holly	<i>Taxus cuspidate</i>	Japanese Yew
<i>Ilex crenata</i>	Japanese Holly		
<i>Ilex opaca</i>	American Holly		
<i>Ilex pernyi</i>	Holly	<i>Taxus media</i>	Yew, Var.
<i>Iris. Spp</i>	Iris species	<i>Thuja occidentalis</i>	American Arborvitae
<i>Jasminium nudiflorum</i>	Winter Jasmine	<i>Trachelospermum asiaticum</i>	Star Jasmine
<i>Juglans</i> sp.	Walnut*	<i>Tsuga canadensis</i>	Canada Hemlock
<i>Juniperus chinensis</i>	Chinese Juniper	<i>Tulipa</i> spp.	Tulip species
<i>Juniperus conferta</i>	Shore Juniper	<i>Viburnum japonicum</i>	Japanese Viburnum
<i>Juniperus horizontalis</i>	Creeping Juniper	<i>Viburnum plicatum</i>	Japanese Snowball
<i>Justica brandegeana</i>	Shrimp Plant	<i>Viburnum trilobium</i>	Cranberry Bush
<i>Lagerstromia indica</i>	Crape Myrtle	<i>Viburnum wrightii</i>	Leatherleaf Viburnum
<i>Ligustrum amurense</i>	Amur Privet	<i>Vinca major</i>	Vinca
<i>Ligustrum japonicum</i>	Japanese Privet	<i>Vinca minor</i>	Dwarf Periwinkle
<i>Ligustrum lucidum</i>	Glossy Privet (Wax-Leaf)	<i>Vitis</i> sp.	Grape*
<i>Liriope muscari</i>	Big Blue Lily Turf	<i>Weigela florida</i>	Old Fashioned Weigela
<i>Lonicera japonica</i>	Japanese Honeysuckle	<i>Yucca aloifolia</i>	Spanish Bayonet
<i>Magnolia</i> spp.	Magnolia species	<i>Yucca filamentosa</i>	Yucca, Adam's Needle

**\*Do not use on food-producing trees, vines, or plants.**  
LESCO Barricade 0.38% Plus Fertilizer controls certain grass and broadleaf weeds before they emerge and provides residual control in established turfgrasses, and ornamentals and other non-crop area. LESCO Barricade 0.38% Plus Fertilizer inhibits shoot and root growth in susceptible weeds. LESCO Barricade 0.38% Plus Fertilizer will not control established weeds. Weed control is most effective when LESCO Barricade 0.38% Plus Fertilizer is activated by at least ½ inch of rainfall or irrigation before weed seeds germinate and within 14 days following application.

## WEEDS CONTROLLED

Barnyardgrass	Goosegrass <sup>3</sup>	Lovegrass	Speedwell, Persian
Bluegrass, Annual ( <i>Poa annua</i> ) <sup>1</sup>	Henbit	Panicum (Texas, Fall, Brown Top)	Sprangletop
Carpetweed	Itchgrass	Pigweed	Spurge, Prostrate
Chickweed, Common, Mouseear (from seed)	Johnsongrass (from seed)	Purslane, Common	Witchgrass
Crabgrass (large, smooth) <sup>2</sup>	Junglerice	Pusley, Florida	Woodsorrel, Yellow (from seed)
Crowfoot grass	Knotweed	Rescuegrass <sup>4</sup>	
Cupgrass, Woolly	Kochia	Shepherd's Purse	
Foxtails, Annual	Lambsquarter, Common	Signalgrass, Broadleaf	

- In areas where *Poa annua* is a winter annual, apply **LESCO Barricade 0.38% Plus Fertilizer** (see rate table) in August or September to established, non-overseeded turf before *Poa annua* seeds germinate. These timings are approximate. Consult State Extension Service for more specific timing for your area.
- Fall Applications for Spring Crabgrass Control in Cool-Season Grasses: In those areas where the ground freezes in the winter, **LESCO Barricade 0.38% Plus Fertilizer** can be applied in the fall after the soil temperature falls below 50F but before the ground freezes. This application will control crabgrass the following spring.
- In many areas, a single application will control goosegrass. However, under heavy goosegrass pressure and/or an extended growing season, make a second application 60-90 days later to ornamentals and lawns (if recommended for your lawn grass species).
- Suppression only.

**EPA REG. NO. 100-1339-10404**  
**EPA EST. NO. 82757-FL-001(S), 82757-OH-001 (M), 82757-MA-002 (H)**

First letter of lot code indicates manufacturing site.

# #097649

# T

## Use Precautions

- Do not apply to putting greens or turfgrass areas where dichondra, colonial bentgrass, velvet bentgrass, or annual bluegrass (*Poa Annua*) are desirable species.
- Do not cut (harvest) treated sod before 90 days after application. To avoid turfgrass injury, do not apply to newly-set sod until the sod has rooted into the second inch of soil and exposed edges have filled in.
- Application of LESCO Barricade 0.38% Plus Fertilizer to turf stressed by drought, low fertility, or pest damage may result in turf injury.
- Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.

**Seeding:** Do not apply to overseeded areas for 60 days after seeding. Or until after the second mowing, whichever is longer. Do not overseed for 4 months after last applications of this product

**Granular Spreader Settings**<sup>1,2,3</sup>

Make and Models	[0.38 lb. ai/A] [2.3 lb./1,000 sq. ft] Base Rate (Treats 21,700 sq. ft.)	[0.5 LB. ai/A] [3.0 lb./1,000 sq. ft] Medium Rate (Treats 16,600 sq. ft.)	[0.75 lb. ai/A] [4.5 lb./1,000 sq. ft] High Rate (Treats 11,000 sq. ft.)
LESCO Calibration Gauge	12	13	16
SCOTTS® R8A	1	1½	K
Cyclone® or Spyker®	3¼	3½	4½
LESCO Pendulum	18	20	30

- Assume 3 mph walking speed.
- It is recommended that you calibrate your spreader before application since manufacturer's settings can vary among the same model of spreader.
- Do not apply more than 1.5 lbs. ai/A per year.

## LANDSCAPE, FIELD GROWN ORNAMENTALS AND NON-CROP AREAS How to Apply

- Apply LESCO Barricade 0.38 Plus Fertilizer as a broadcast, over-the-top application at 3.9-9.0 lbs. product/1,000 sq. ft. Use higher rates when weed pressure was a severe problem in the past.
- Make a single application in fall and/or spring prior to weed emergence. Another application may be made 60-90 days later if necessary. Do not exceed a total of 9.0 lbs. product/1,000 sq. ft. per year.
- After application, water in ½ inch of water within 14 days, either from watering or rain or, incorporate into the top 1-2 inches of soil within 14 days after application.
- On established ornamentals, watering or rainfall soon after application will wash residues off plant foliage and activate LESCO Barricade 0.38 Plus Fertilizer in the soil.
- On newly-transplanted ornamentals, apply after the soil has settled around the transplants.
- For best weed control, apply to soil free of clods, weeds, and debris such as leaves.
- Prior to application, control existing weeds by hand weeding, cultivation, or by the use of an herbicide for emerged weeds.
- Plants under stress (such as heat, drought, or frost damage) may be sensitive to LESCO Barricade 0.38 Plus Fertilizer.

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

**PESTICIDE STORAGE:** Store in a cool, dry place. Keep container closed and away from food, feedstuffs, and domestic water supplies.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Non-refillable container. Do not reuse or refill this container. Completely empty bag into application equipment. Offer for recycling if available or dispose of bag in a sanitary landfill or by other procedures approved by state and local authorities.

## CONDITION OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.  
The Directions for Use of this product must be followed carefully and completely. It is impossible to eliminate all risks inherently associated with the use of this product. To the extent consistent with applicable law, Buyer and/or User assume all risks of ineffectiveness or other unintended consequences or damages that may result from conditions outside or beyond the control of LESCO, Inc. including but not limited to, such factors as manner of use or application, weather or weather conditions outside the range considered normal at the application site or for the time period in which the product is applied, the presence of other materials, incompatible products, or other influencing factors which are beyond the control of LESCO, Inc.. To the extent consistent with applicable law, all such risks shall be assumed by Buyer and/or User, and Buyer and/or User agrees to hold LESCO, Inc. harmless for any claims relating to such factors.  
LESCO, Inc. warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with the Directions for Use under normal use conditions. To the extent consistent with applicable law, this warranty does not extend to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of LESCO, Inc. and Buyer and/or User assume the risk of any such use. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, LESCO, INC. MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.**  
**TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE BUYER AND/OR USER AND THE EXCLUSIVE LIABILITY OF LESCO, INC. FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF LESCO, INC. REPLACEMENT OF THE PRODUCT, OR IF NOT ACQUIRED BY PURCHASE, REPLACEMENT OF SUCH QUANTITY. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, IN NO EVENT SHALL LESCO, Inc., BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES.**  
LESCO, Inc. offers this product, and Buyer and/or User accepts it, subject to the foregoing Conditions of Sale and Limitation of Warranty and Liability, which may not be modified except by written agreement signed by a duly authorized representative of LESCO, Inc.

Information concerning the raw materials composing this product can be obtained by writing to: LESCO, Inc., Attn: RA Dept., 1385 East 36<sup>th</sup> Street, Cleveland, Ohio 44114-4114, referring to the item number found on this bag.  
Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.htm>  
LESCO is a registered trademark and the sweeping design is a trademark of LESCO Technologies, LLC. Barricade is a trademark of a Syngenta Group Company. SCOTTS is a registered trademark of The SCOTT Company. Cyclone and Spyker are registered trademarks of Spyker Spreaders, LLC.  
G:\REGUL\WP\Erview\Aprvrd-Lbl Word\097649.doc  
Created: 11/30/09 (090810) Rev.: 8/16/12 VT  
**Net Weight: 50 lb (22.7 kg)**

## F1560

Distributed by: **LESCO, Inc.**  
1385 East 36<sup>th</sup> Street  
Cleveland, OH 44114-4114